

Natural colorants: Pigment stability and extraction yield enhancement via utilization of appropriate pretreatment and extraction methods

Luxsika Ngamwonglumert, Sakamon Devahastin & Naphaporn Chiewchan

To cite this article: Luxsika Ngamwonglumert, Sakamon Devahastin & Naphaporn Chiewchan (2017) Natural colorants: Pigment stability and extraction yield enhancement via utilization of appropriate pretreatment and extraction methods, *Critical Reviews in Food Science and Nutrition*, 57:15, 3243-3259, DOI: [10.1080/10408398.2015.1109498](https://doi.org/10.1080/10408398.2015.1109498)

To link to this article: <https://doi.org/10.1080/10408398.2015.1109498>



Accepted author version posted online: 30 Oct 2015.
Published online: 23 May 2017.



Submit your article to this journal [↗](#)



Article views: 1447



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 28 View citing articles [↗](#)



Natural colorants: Pigment stability and extraction yield enhancement via utilization of appropriate pretreatment and extraction methods

Luxsika Ngamwonglumlert, Sakamon Devahastin , and Naphaporn Chiewchan

Advanced Food Processing Research Laboratory, Department of Food Engineering, Faculty of Engineering, King Mongkut's University of Technology Thonburi, Tungkrui, Bangkok, Thailand

ABSTRACT

Natural colorants from plant-based materials have gained increasing popularity due to health consciousness of consumers. Among the many steps involved in the production of natural colorants, pigment extraction is one of the most important. Soxhlet extraction, maceration, and hydrodistillation are conventional methods that have been widely used in industry and laboratory for such a purpose. Recently, various non-conventional methods, such as supercritical fluid extraction, pressurized liquid extraction, microwave-assisted extraction, ultrasound-assisted extraction, pulsed-electric field extraction, and enzyme-assisted extraction have emerged as alternatives to conventional methods due to the advantages of the former in terms of smaller solvent consumption, shorter extraction time, and more environment-friendliness. Prior to the extraction step, pretreatment of plant materials to enhance the stability of natural pigments is another important step that must be carefully taken care of. In this paper, a comprehensive review of appropriate pretreatment and extraction methods for chlorophylls, carotenoids, betalains, and anthocyanins, which are major classes of plant pigments, is provided by using pigment stability and extraction yield as assessment criteria.

KEYWORDS

Anthocyanin; betalains; carotenoids; chlorophylls; natural products; non-conventional extraction

Introduction

Among many characteristics of food, color is one of the most important one, as it can lead to a good first impression of a product. Unfortunately, most food processing operations, especially those involving the use of heat, generally causes alteration, degradation, or even loss of food color. For this reason, a wide array of food colorants (i.e. pigments and dyes) has been added into food to enhance or intensify its original color. Colorants can also be added to ensure color uniformity or to obtain the best food appearance, or even to provide color to otherwise uncolored food (Delgado-Vargas and Paredes-López, 2003).

Most food colorants that are in use today are derived from minerals or synthetic processes. Seven types of synthetic pigments, which are Brilliant Blue FCF (FD&C Blue No. 1), Indigotine (FD&C Blue No. 2), Fast Green (FD&C Green No. 3), Erythrosine (FD&C Red No. 3), Allura Red AC (FD&C Red No. 40), Tartrazine (FD&C Yellow No. 5), and Sunset Yellow FCF (FD&C Yellow No. 6), are permitted in foods under the Food and Drug Administration (FDA) regulations (Harp and Barrows, 2015). Although synthetic colorants possess higher stability, more diverse hue, and vibrant color, their consumption has been reported to be related to many adverse health effects such as attention problems, hyperactivity, irritability, sleep disorders, and aggressiveness in children (Masone and Chanforan, 2015). Due to health consciousness of modern consumers, colorants produced from natural sources (natural colorants) are an attractive alternative and have gained increasing

popularity. Chlorophylls, carotenoids, betalains, and anthocyanins are four important classes of natural pigments that are abundant in nature and their extracts have been approved as food additives by FDA and Codex (Francis, 1996). Nevertheless, less stability and limited diversity in terms of color shade are still problems of natural colorants (Delgado-Vargas and Paredes-López, 2003).

In general, to produce a natural colorant, extraction process is first required to extract crude pigment from a starting material, which most of the time is a plant material. Extraction of a plant material can be done by various extraction techniques. Normally, conventional methods such as Soxhlet extraction are used. More recently, however, many non-conventional methods, such as supercritical fluid extraction (SFE), pressurized liquid extraction (PLE), microwave-assisted extraction (MAE), ultrasound-assisted extraction (UAE), pulsed-electric field (PEF) extraction, and enzyme-assisted extraction (EAE), have been proposed due to their enhanced extraction efficiency and environment-friendliness (Azmir et al., 2013). A suitable extraction technique helps to increase extraction yield and prevent the degradation of extracted pigments, leading to an ability to produce natural colorants of higher quality.

A number of researchers have also proposed the use of various pretreatment methods prior to extraction to increase extraction yield and/or enhance the stability of pigments. Thermal pretreatments (e.g., steam and hot water blanching) and chemical soaking are among the most popular methods that

can be used to pre-treat a plant material. Thermal treatments are mainly used to inactivate deleterious enzymes, which cause pigment degradation; these treatments also help induce changes in the physical structure of a plant material, resulting in an enhanced extraction yield (Heaton and Marangoni, 1996; Stamatopoulos et al., 2012). Enhancement of pigment stability by using chemical solutions (e.g., organic acid solutions, alkaline solutions, and ionic solutions) is also well recognized (Maharaj and Sankat, 1996; Reynoso et al., 1997; Bąkowska et al., 2003; Hiranvarachat et al., 2011). However, selecting a proper pretreatment method for each pigment group is of paramount importance and must always be done with great care.

This paper aims to provide a comprehensive review of appropriate pretreatment and extraction methods for each group of four important pigments, namely, chlorophylls, carotenoids, betalains, and anthocyanins, by using stability and extraction yield of pigments as assessment criteria.

Important classes of natural pigments and their stability

Two systems have been used to classify natural pigments. The first one is based on structural affinities; natural pigments are divided into six major structural classes, which are tetrapyrroles, tetra-terpenoids, quinones, *N*-heterocyclic, *O*-heterocyclic, and metallo-proteins (Hendry, 1996). This system has an advantage of brevity, but biosynthetic compounds, which are quite unrelated, are grouped together. The other classification system is based on natural occurrence of pigments (Hendry, 1996), and is adopted in the present paper. According to this system of classification, pigments in plant materials are classified into chlorophylls, carotenoids, betalains, and anthocyanins.

Chlorophylls

Chlorophylls are oil-soluble, and the most widely distributed pigments responsible for the characteristic green color of plants. The structure of chlorophylls is porphyrins or closed ring tetrapyrroles chelated with a centrally located magnesium atom (Marquez and Sinnecker, 2008), as shown in Fig. 1. There are five classes of chlorophylls that have been found in plants and photosynthetic organisms. However, the major chlorophylls found in foods are chlorophyll *a* and chlorophyll *b*. These chlorophylls are different at position 7, where chlorophyll *a* is composed of $-\text{CH}_3$ and chlorophyll *b* is composed of $-\text{CHO}$. This difference leads to difference in color; chlorophyll *a* appears blue-green, while chlorophyll *b* possesses yellow-green color (Steet and Tong, 1996).

Chlorophylls are highly sensitive to heat, light, oxygen, acids, and enzymes, leading to their easy degradation and color change (Özkan and Bilek, 2015). Acids and Mg-dechelataase, which is an enzyme found in algae and plants, are the main causes of conversion of chlorophyll's color from green to olive brown (Marquez and Sinnecker, 2008). These changes occur due to the loss of central magnesium atom in the chlorophyll's structure, with the replacement by hydrogen ions, leading to the transformation of chlorophyll's structure from native chlorophyll to pheophytin, which exhibits olive brown color




Figure 1. Basic structure of chlorophylls pigment (Marquez and Sinnecker, 2008).

(Delgado-Vargas and Paredes-López, 2003). Other enzymes, such as chlorophyllase and oxidative enzymes (e.g., lipoxygenase, chlorophyll oxidase, and peroxidase), are also related to chlorophyll's degradation and cause change in colors (Marquez and Sinnecker, 2008). Thus, inhibition of enzyme activity and avoidance of acidic condition must be exercised to prevent changes in chlorophyll's color.

Heat is another important factor that indirectly causes conversion in chlorophyll's structure, which subsequently leads to color changes. Many researchers have investigated the effect of thermal processes on change in chlorophyll's color. For example, Lau et al. (2000) studied the color change of green asparagus during thermal treatment and found that the color of the sample changed to olive color due to degradation of chlorophylls by heat. In fact, heat simply induces breaking up or disruption of plant cell walls; intracellular acids are thus released from the cells, resulting in the change of pH to an acidic condition (Fellows, 2000; Marquez and Sinnecker, 2008). For these reasons, the structure of native chlorophylls changes to pheophytin, and hence the color changes from green to olive brown as mentioned earlier.

Carotenoids

Carotenoids are lipid-soluble pigments responsible for many of the brilliant red, orange, and yellow colors of fruits, vegetables, and flowers (Ötles and Çagindi, 2008). Based on their structure, carotenoids can be divided into two groups, which are carotenes and xanthophylls (Fig. 2). Carotenes are constituted by carbon and hydrogen, whereas xanthophylls are constituted by carbon, hydrogen, and oxygen (Britton, 1996). Carotenoids can be divided based on their functionality into primary and




Figure 2. Structures of major carotenenes (I) and xanthophylls (II) (Ötles and Çagindi, 2008).

secondary carotenoids. Primary carotenoids (e.g., β -carotene, lutein, zeaxanthin, and antheraxanthin) are the pigments required for photosynthetic process, whereas secondary carotenoids (e.g., α -carotene, lycopene, astaxanthin, and canthaxanthin) are the pigments that are not directly related with plant survival (Delgado-Vargas et al., 2000).

Degradation of carotenoids is mainly caused by the reactions of oxidation and isomerization, which lead to a decrease in the redness and yellowness of plant material (Provesi and Amante, 2015). There are several factors affecting the occurrence of oxidation and isomerization of carotenoids. Oxygen naturally causes oxidation of carotenoids; the reaction can also be stimulated by light, heat, peroxide, metal ions, and enzymes (Boon et al., 2010). Most carotenoids in plants are *trans*-isomers, and the isomerization of *trans*- to *cis*-isomers occurs during food processing (Khoo et al., 2011). Heat and light as well as acids are the main factors that can promote isomerization of carotenoids from *trans*- to *cis*-isomers (Provesi and Amante, 2015). However, isomerization of carotenoids leads to slightly reduced biological activities and color saturation, whereas oxidation leads to complete loss in the activities and color of carotenoids (Takyi, 2001). Several precautions (e.g., processing under low pressure, storage in dark under low temperature and low oxygen conditions) have therefore been proposed to prevent oxidation reaction (Ötles and Çagindi, 2008; Khoo et al., 2011).

Betalains

Betalains are *N*-heterocyclic water-soluble pigments, which exhibit red–purple or yellow color depending on the pigment structure (Delgado-Vargas and Paredes-López, 2003). These pigments are found in many fruits, vegetables, and flowers (e.g., prickly pear, red beetroot, aizoaceae). Betalains can be classified into two groups based on their structure, which are betaxanthins (yellow pigment) and betacyanins (red–purple pigment) as shown in Fig. 3 (Chandrasekhar et al., 2015).

Instability of betalains to light, heat, alkaline, oxygen, and metal ions restricts them from extensive applications in food (Khan and Giridhar, 2014). Among various factors, heat is the most critical factor affecting the degradation of betalains (Reshmi et al., 2012), while other factors affect the heat sensitivity of betalains. Heat mostly induces oxidation (i.e., dehydrogenation in the presence of oxygen), aldimine bond hydrolysis, and decarboxylation of betalains, which results in color change to orange-yellow (Gonçalves et al., 2013). Since betalains are susceptible to heat, it is most important to cool down immediately the treated plant material after thermal treatment (Henry, 1996).

Besides heat, stability of betalains is also related to pH. These pigments are most stable in a pH range of 3 to 7. Betacyanins are more resistant to acidic condition, while betaxanthins are




Figure 3. Structures of betaxanthins (I) and betacyanins (II) (adapted from Herbach et al., 2006a).

most stable at neutral pH (Stintzing and Carle, 2008). The color of betalains does not change within the pH range of 3 to 7. When the pH is below 3, the structure of betalains is converted from red anionic to violet cationic, resulting in color change from red to blue–violet shade (Henry, 1996). On the other hand, alkaline condition (pH higher than 7) brings about aldimine bond hydrolysis, which results in rapid degradation of betalains into betalamic acid (BA) and cyclodopa-5-*O*-glucoside (CDG) and the color change to yellow-brown (Henry, 1996).

Metal ions (e.g., Sn^{2+} , Al^{3+} , Ni^{2+} , Cr^{2+} , Fe^{2+} , Fe^{3+} , and Cu^{2+}), which may contaminate a plant (from soil accompanying the plant), can accelerate oxidation of betalains, resulting in loss in colors (Stintzing and Carle, 2008). However, contamination of metal ions can be minimized by washing. Light does not significantly cause degradation of this group of pigments (Ötles and Çagindi, 2008).


Anthocyanins

Anthocyanins are water-soluble pigments found in many flowers, fruits, and leaves (e.g., butterfly pea, red onion, grape, and purple corn) with various shades (orange, red, or blue) (Delgado-Vargas and Paredes-López, 2003). The basic structure of anthocyanins is glycosides of anthocyanidins, which is known as flavylium cation (Mercadante and Bobbio, 2008). The term anthocyanidins is referred to anthocyanins without sugar molecules. The most abundant anthocyanins in nature are based on six anthocyanidins (Wu and Prior, 2005), which are pelargonidin (plg), cyanidin (cyd), delphinidin (dpd), penodan (pnd), petunidin (ptd), and malvidin (mvd), as shown in Fig. 4.

Differences in the number of hydroxyl and methoxyl groups in the structure affect the color variation of pigments. Pigments that have a larger number of hydroxyl groups exhibit more bluish shade, while those having a larger number of methoxyl groups exhibit more redness (Delgado-Vargas and Paredes-López, 2003).

The color stability of anthocyanins is influenced not only by structural features but also by pH, temperature, light, presence of co-pigments, enzymes, oxygen, and sugars (Rodríguez-Saona et al., 1999). Change in pH causes reversible transformation in anthocyanins structure and significantly affects shift in the color of anthocyanins. In an aqueous solution, the molecular species of anthocyanins, which are flavylium cation, carbinol pseudobase, quinoidal base, and chalcone, are in equilibrium. Flavylium cation, which exhibits red color, dominates at pH 1, but when the pH increases to 4 or 5, the concentration of carbinol pseudobase, which is colorless, increases at an expense of flavylium cation. At pH 6, hydroxyl groups present in the structure and the concentration of quinoidal base increase, yielding unstable blue or violet color. Quinoidal base continuously increases until the pH rises to 8; quinoidal base would then transform into chalcone, which exhibits yellow color (Delgado-Vargas and Paredes-López, 2003; Ananga et al., 2013). Structural transformation and color change of anthocyanins with pH are shown in Fig. 5.

Heat is well known as an important factor affecting the color and stability of natural pigments. However, anthocyanins are more stable to heat than the aforementioned pigments. Degradation of anthocyanins occurs when temperature approaches 100°C or even higher (Jackman and Smith, 1996). The first step of thermal degradation involves the formation of colorless



Anthocyanidin	Substitution pattern			Color
	R ₁	R ₂	R ₃	
Pelargonidin (plg)	H	OH	H	Orange
Cyanidin (cyd)	OH	OH	H	Magenta, Crimson
Delphinidin (dpd)	OH	OH	OH	Purple, mauve, blue
Peonidin (pnd)	OCH ₃	OH	H	Magenta
Petunidin (ptd)	OCH ₃	OH	OH	Purple
Malvidin (mvd)	OCH ₃	OH	OCH ₃	Purple

Figure 4. Structures of six important anthocyanidins in nature (adapted from Delgado-Vargas and Paredes-López, 2003; Mercadante and Bobbio, 2008).

carbinol pseudobase and the subsequent opening of pyrylium ring to form chalcone, before transforming into a brown degradation product (Perera and Baldwin, 2001).

The color stability of each pigment is briefly summarized in Table 1.

Stability enhancement via pretreatments

Due to the instability of natural pigments, various pretreatment methods have been suggested to enhance stability and modify the structure of plant materials. Pretreatment methods may be divided into three groups, which are physical, chemical, and biological pretreatments. Chemical pretreatments (e.g., acid and alkaline treatments) are proposed to enhance the stability of pigments. On the other hand, the main purpose of physical (e.g., microwave heating, PEF extraction, and ultrasonic treatments) and biological (e.g., enzymatic treatment) pretreatments is plant structural modification to enhance subsequent extraction. The use of physical and biological pretreatments in combination with extraction will be discussed in the later section. Since the stability depends on the type of pigments, a suitable method of pretreatment that should be used to increase the stability also depends on pigment-types.

Stability enhancement of chlorophylls

Enzymes and acids are the major factors affecting the degradation of chlorophylls as mentioned in the section “Chlorophylls.” Thus, enhancing the stability of chlorophyll pigments may be done by inactivating the enzymes and/or preventing an

acidic condition. Many pretreatment methods have been proposed for such purposes. Blanching is a short-time thermal treatment, which is usually performed by subjecting a sample to hot water or steam, with the aim to inactivate enzymes involved in the change of color of plant materials (Bahçeci et al., 2005). Most enzymes normally suffer decreased activity when being heated at above 80°C (von Elbe and Schwartz, 1996). Since peroxidase is one of the highly thermal resistant enzymes and its activity is easy to measure, it is generally used as an indicator of adequate blanching (Bahçeci et al., 2005; Viña et al., 2007). Olivera et al. (2008), for example, reported that blanched Brussels sprouts exhibited more greenness than the unbalanced sample after storage at –18°C for eight months. However, after eight months of frozen storage, the greenness values of both blanched and unblanched samples decreased. Blanching nevertheless leads to changes in the cellular structure of plant, sometimes resulting in undesirable softening of plant tissues (Fellows, 2000). In addition, blanching cannot solve the problem of acidic condition that may occur in further processing steps.

Alkaline treatment is proposed to prevent acidic condition. Alkalinizing agents (e.g., calcium hydroxide, magnesium hydroxide, potassium hydroxide, sodium hydroxide, and potassium bicarbonate) in combination with blanching are usually used to enhance the stability of green color. Alkalinizing agents raise the pH of a blanched sample, thus enabling the retention of green color of chlorophylls (Negi and Roy, 2001; Ahmed et al., 2013). Maharaj and Sankat (1996) reported that blanching dasheen leaves in alkaline solution (0.06% magnesium carbonate solution) at 100°C for 10 s prior to drying at 60°C




Figure 5. Changes in the structure and color of anthocyanins at different pH values (adapted from Ananga et al., 2013).

resulted in better retention of green color of the dried product when compared with the products obtained from unblanched and even steam-blanched samples. Unfortunately, commercial application of alkaline treatment is still not too effective, as alkalinizing agents do not have the ability to neutralize interior tissue acids over a long period of time, leading, for instance, to the change in the color of canned green peas after less than two months of storage at room temperature (von Elbe and Schwartz, 1996).

Complexation of chlorophylls into metal complexes of chlorophyll derivatives (e.g., copper pyropheophytin and zinc pyropheophytin) has been proposed to alleviate the aforementioned problems. Chlorophylls can be transformed into metallo-chlorophyll derivatives, which exhibit green color similar to native chlorophylls but are more stable to acids and heat (Marquez

and Sinnecker, 2008). Although both copper and zinc ions can be utilized for metallo-chlorophyll derivatives formation, zinc ions are of greater interest due to the toxic nature of copper ions (Delgado-Vargas and Paredes-López, 2003).

To create metallo-chlorophyll derivatives, chlorophylls are first acidified to change the structure of chlorophylls to pheophytin prior to blanching in a medium containing metal ions, such as zinc or copper ions, to form metallo-chlorophyll derivatives (Tonucci and von Elbe, 1992) as illustrated in Fig. 6. Many researchers have indeed investigated the thermal stability of these metallo-chlorophyll derivatives and reported improved results. For example, Ngo and Zhao (2005) reported that blanching green pear chunks at 94°C in 2600-ppm Zn²⁺ solution for 13 min could preserve the green color of samples during sterilization process (94°C, 20 min). The color of the

Table 1. Color stability of chlorophylls, carotenoids, betalains, and anthocyanins to heat, light, oxygen, and pH.

Pigment	Typical source	Stability			
		Heat	Light	Oxygen	pH change
Chlorophylls	Leaves	Moderate	High	High	Low
Carotenoids	Leaves	Moderate	Low	Low	High
Betalains	Beetroot	Moderate to low	High	High	Moderate
Anthocyanins	Fruits	High to moderate	High	High	Low




Figure 6. Chemical reactions for creating zinc-chlorophylls derivative (adapted from Socaciu, 2008).

sample did not change for 19 weeks at 38°C under intensive illumination or for at least 35 weeks at 10°C.

Stability enhancement of carotenoids

Degradation of carotenoids with their highly conjugated double-bond structure is mainly due to oxidation reaction (von Elbe and Schwartz, 1996). Inactivation of oxidative enzymes is among the possible means of solving the problem of carotenoids oxidation. Hot water or steam blanching is a simple pretreatment method that can be used to inactivate enzymes such as lipoxygenase, which catalyzes oxidative decomposition of carotenoids (von Elbe and Schwartz, 1996). However, although blanching has a benefit of enzyme inactivation, many investigators have reported that blanching in some cases led to the degradation of carotenoid pigments (Hiranvarachat et al., 2011). Marx et al. (2003) and Lavelli et al. (2007) who studied the effect of blanching on the carotenoids degradation of carrot juice and dehydrated carrot, respectively, reported that carrot juice and freeze-dried carrot produced from blanched carrot suffered more extensive degradation of carotenoids than those produced from unblanched carrot; the degradation led to the fading color of products. This observation might be caused by the change in carotenoid structure from *trans*- to *cis*-isomers due to heat upon blanching.

Chemical pretreatments are another alternative method to prevent the oxidation of carotenoids. Antioxidant agents (e.g., butylated hydroxyanisole, pyrogallol, and citric acid) are proposed to retard pigment degradation by oxidation (Ötles and Çagindi, 2008). Hackett et al. (2004) reported that addition of antioxidants (α -tocopherol or butylated hydroxytoluene) helped decrease the degradation rate of lycopene in tomato heated at 50°C. Hiranvarachat et al. (2011) showed that β -carotene content of carrot soaked in citric acid (pH 4.0–5.0) tended to be unchanged during drying (hot air drying), while β -carotene content of untreated carrot decreased continuously. However, adding acid-based antioxidants leads to a decrease in the pH value of a sample; when the pH of a sample becomes less than 3.0, carotenoids would start to degrade. Note that carotenoids would significant degrade at pH below 3.0, whereas these become more stable at pH between 4.0 and 6.0. Under acidic condition (pH < 3.0), carotenoids are protonated, resulting in the isomerization of their structure from *trans*- to *cis*-isomers (Qian et al., 2012). Thus, the pH value of a sample should be

carefully controlled when acid-based antioxidants, such as organic acids, are used.

Stability enhancement of betalains

To improve the stability of betalains, blanching is usually required to inactivate the betacyanin decoloring enzyme, which leads to color fading (Delgado-Vargas and Paredes-López, 2003). Nevertheless, heat is the most crucial factor affecting the stability of betalains. As mentioned in the section “Betalains,” the color loss due to the degradation of betalains is mainly caused by heat and alkalinity, which induce aldimine bond hydrolysis, leading to the separation of betalains structure into BA and CDG (Henry, 1996).

Betalains can be regenerated after thermal processing via the use of acids (Herbach et al., 2006b). Various organic acids (e.g., ascorbic acid, isoascorbic acid, citric acid, and gluconic acid) have been proposed to aid regeneration and retain stability during processing and storage of betalains. Bilyk et al. (1981) noted that addition of isoascorbic acid into a betacyanins solution either before or after heating (at 100°C) helped to regenerate betacyanins. The regenerated betacyanins were restored almost completely after 24 h (storage in dark at 25°C). On the other hand, the degraded betaxanthins could not be regenerated by the use of isoascorbic acid due to its instability in acidic condition (Stintzing and Carle, 2008). Then organic acid is a suitable pretreatment for enhancing the stability of red colorant from betacyanins but not for yellow colorant from betaxanthins.

The quantity of regenerated betacyanins is affected by pH, storage temperature, type of additives, and the presence or absence of oxygen (Bilyk and Howard, 1982). Han et al. (1998) studied the regeneration of degraded red beet betacyanins in the presence of different types of acid-based additives, which were antioxidants (ascorbic acid, isoascorbic acid, and gluconic acid), organic acids (acetic, citric, lactic, gallic, and gluconic acids), and inorganic acids (metaphosphoric acid, phosphoric acid, and Na_2 EDTA). Acid-based additives helped to regenerate red beet pigments that were degraded by heating at 100°C for five minutes. The degraded pigments were restored under 10°C for 10 min. The effect of pH on pigment regeneration was also evaluated, and it was found that the addition of ascorbic acid and isoascorbic acid resulted in the highest retention of pigments (98.3%) for the sample controlled at pH 3.0. Addition

of gluconic acid and metaphosphoric acid also resulted in the higher retention of pigments, with the retention of 81.7% and 75.4%, respectively for the sample controlled at pH 6.8.

Besides the ability to regenerate the pigment after heating, ascorbic acid and isoascorbic acid could also help stabilize betacyanins during food processing and storage. Reynoso et al. (1997) reported that garambullo juice that was added with 0.1% ascorbic acid exhibited higher redness than the untreated sample after sterilization at 121°C for 15 min. The presence of the acid in red beet juice and garambullo juice (adjusted pH 5.5) also resulted in an increase in the stability of betalains during storage at 25°C for five days. Moreover, ascorbic acid could improve the stability of betalains when metal ions (chromium and iron) are present, since the acid could act as a chelating agent (Pokorny, 2007).

Stability enhancement of anthocyanins

Anthocyanins are highly sensitive to change in pH, which results in shift of color. Using anthocyanins as a natural colorant is suggested at low pH (pH < 4.0). At pH below 4, anthocyanins primarily are in the form of flavylium cation, which is more stable than other structures (Tan et al., 2014). Cevallos-Casals and Cisneros-Zevallos (2004), for instance, reported that at a pH in the range of 0.9 to 4.0, the colorant samples from red sweet potato and purple carrot showed high stability during storage at 20°C for 134 days; hue angle of the red sweet potato colorant at pH 4.0 still exhibited red-violet color after storage. In contrast, the color of the samples that were controlled at pH above 4.0 shifted from purple-blue to brown and yellow, and after 134 days the color of all the samples was predominated by the yellow-colored chalcone.

Besides a proper pH adjustment, the source of anthocyanins must be considered to obtain the most stable colorant. Different sources of plant materials contain different anthocyanin structures, which affect their stability. Anthocyanins from red cabbage, black carrot, red radish, and red sweet potato are reported to be more stable to heat and pH change than anthocyanins from other sources due to their acylation of structure (Bąkowska-Barczak, 2005). Acylation of anthocyanin molecules can enhance stability through intramolecular co-pigmentation (Rodríguez-Saona et al., 1999). Higher stability of acylated anthocyanins is attributed to the stacking of acyl group with the pyrylium ring of flavylium cation, thus preventing the nucleophile attack of water and subsequent formation of chalcone (Bąkowska-Barczak, 2005). Many researchers have indeed demonstrated the stability of acylated anthocyanins in comparison with that of non-acylated anthocyanins. For example, Cevallos-Casals and Cisneros-Zevallos (2004), who studied the stability of anthocyanins from red sweet potato, purple carrot, purple corn, and red grape, reported that anthocyanins from red sweet potato and purple carrot, which mainly consist of acylated anthocyanins, were more stable than those from purple corn and red grape, which mainly consist of non-acylated anthocyanins. Anthocyanins extracted from red sweet potato exhibited longer half-life than those from purple carrot, purple corn, and red grape when heated at 98°C and controlled at pH 3.0.

Intercellular co-pigmentation has been suggested to improve the stability of anthocyanins. Intercellular co-pigmentation is

defined as interactions between anthocyanin molecules and other molecules, e.g., flavonoids, alkaloids, amino acids, organic acids, metals, and other anthocyanins (Castañeda-Ovando et al., 2008). The basic role of intercellular co-pigmentation is the same as that of intracellular co-pigmentation, which is to protect flavylium cation from the nucleophile attack of water molecules (Mazza and Brouillard, 1990; Castañeda-Ovando et al., 2008). Gauche et al. (2010) reported that addition of organic acids (caffeic, ferulic, gallic, and tannic acids) into an anthocyanins solution could retard color change when the pH of the solution increased from 4.0 to 6.0, whereas the solution without organic acids became colorless. Moreover, the half-life of anthocyanins solutions with added organic acids was longer than that of the untreated sample during storage at 28°C; addition of tannic acid resulted in the longest half-life of the sample maintained at pH 1.0.

Although intercellular co-pigmentation can help increase the stability of anthocyanins, this reaction induces an increase in the absorbance (hyperchromic effect: ΔA) and wavelength of the maximum absorbance (bathochromic shift: $\Delta \lambda$) of pigment. Bathochromic shift results in color change from red to red-orange or blue (Bąkowska et al., 2003). Addition of tannic acid (1:1 w/v) to an anthocyanins solution from Isabel grapes (*Vitis labrusca* L.) controlled at pH 3.0 and 4.0, for example, led to an increase in the absorbance and wavelength of the solution, resulting in bluer and brighter color of the treated sample (Bordignon-Luiz et al., 2007). Gauche et al. (2010) also illustrated that co-pigmentation of anthocyanins with tannic, gallic, caffeic, and ferulic acids resulted in hyperchromic effect and bathochromic shift at every pH value (1.0, 2.0, 3.0, 3.3, 3.5, 3.7, 4.0, 4.5) studied. Exception was noted in the case of caffeic acid and ferulic acid at pH of 1.0 and 2.0, where only bathochromic shift was observed. At pH of 1.0 and 2.0, these hydroxycinnamic acids induced the formation of pyranoanthocyanin, which is a anthocyanin-derived pigment, hiding the hyperchromic effect of co-pigmentation (Gómez-Miguez et al., 2006).

Since only a few natural blue-tone colorants are commercially available, production of blue natural colorants from anthocyanins sources is of interest. Blue hue colorants can be produced by the complexation of metal-anthocyanins. However, metal-anthocyanin complexes (blue colorant) are reported to be stable only in the vacuolar matrix (Buchweitz et al., 2013) and would rapidly precipitate in an aqueous solution (Buchweitz et al., 2012). Addition of some polysaccharides or gelatin to form a gel structure is proposed to enhance the stability of metal-anthocyanin complexes by preventing the precipitation of complexes. Buchweitz et al. (2013) noted that gelatin and blend agar-agar with amidated pectin could improve the stability of blue color of ferric anthocyanin chelates during storage at both 20°C in dark and 25°C under illumination. The most stable sample was obtained by using gelatin as a gel matrix; storage should be done at 20°C in dark.

Although the stability of anthocyanins can be enhanced by both intra- and intercellular co-pigmentation as well as preparation of gel matrix as mentioned earlier, controlling the pH value of a food product should be done to prevent shift in color. Storage should be done in dark at cool condition (Bordignon-Luiz et al., 2007).

Examples of stability enhancement of chlorophylls, carotenoids, betalains, and anthocyanins via pretreatments are given in Table 2.

Table 2. Pigment stability enhancement via pretreatments.

Pigment	Source	Pretreatment method	Result	Reference
Chlorophylls	Rocket (<i>Eruca sativa</i>)	Combination of blanching and acid/alkaline treatment	<ul style="list-style-type: none"> Sample blanched in 0.1% NaOH exhibited maximum greenness, while blanching sample in HCl yielded drop in green color. 	Ahmed et al., 2013
	<i>Stevia rebaudiana</i> leaves	Copper treatment	<ul style="list-style-type: none"> Cu-chlorophylls suffered less color loss than Mg-chlorophylls after storage at 60°C for 20 days. Color loss was 16% and 36% for Cu-chlorophyll and Mg-chlorophyll complexes, respectively. 	Bobbio and Guedes, 1990
	Green pear	Blanching and zinc treatment	<ul style="list-style-type: none"> Green pigments of control sample and sample blanched in water were destroyed after heating at 94°C for 12 min. Blanching and zinc treatment helped retain the green pigment of sample after thermal processing. 	Ngo and Zhao, 2005
Carotenoids	Pumpkin (<i>Cucurbita moschata</i> , Duchesne ex Poiret)	Adjusting pH by addition of ascorbic acid and potassium sorbate	<ul style="list-style-type: none"> Addition of ascorbic acid and potassium sorbate to pH 4.0 minimized loss of redness and yellowness of sample after storage at 25°C for 6 weeks. 	Gliemmo et al., 2009
	Carrot	Blanching and acid treatment	<ul style="list-style-type: none"> β-carotene content of sample treated by blanching in citric acid (targeted pH 4–5) tended to be unchanged during thermal processing, whereas β-carotene content of untreated sample decreased continuously. 	Hiranvarachath et al., 2011
Betalains	<i>Rivina humilis</i> L. berry juice	Addition of ascorbic acid	<ul style="list-style-type: none"> Addition of ascorbic acid (0.25 g/100 mL) helped retain betacyanins at 93% and 78% after heating at 90°C for 3 min every 24 h for 6 consecutive days and after heating at 90°C for 24 min, respectively. 	Khan and Giridhar, 2014
Anthocyanins	Blood orange juice	Addition of ascorbic acid	<ul style="list-style-type: none"> Total carotenoids content of sample added with ascorbic acid (30 mg/100 mL) decreased only 2.8%, while that of untreated sample decreased up to 6.6% after storage at 4.5°C for 7 weeks. 	Choi et al., 2002
	Honeysuckle (<i>Lonicera kamtschatica</i>)	Co-pigmentation by using QSA, NaMSA, rutin, quercetin, chlorogenic acid, tannic acid, and flavones	<ul style="list-style-type: none"> Color of co-pigment anthocyanin complexes slowly changed during heating at 180°C for 1 h at pH of 2.5, 3.5, and 4.5 	Bąkowska et al., 2003
	Cabernet Sauvignon grape	Adjusting pH by 0.1-M citric acid–sodium citrate and co-pigmentation by using caffeic acid	<ul style="list-style-type: none"> Co-pigmentation by adjusting to pH 4 yielded retention of color by 96% after storage at 4°C for 72 h, while control sample had only 90% retention of color. Sample extracts added with caffeic acid and adjusted to pH 4 had a half-life of anthocyanins up to 291 days when stored at 4°C in dark, while control sample had a half-life of only 96 days. 	Gris et al., 2007

Effects of extraction methods on pigment extractability

Extraction of pigments from plant materials can be performed by various techniques. Conventional methods are often used to extract crude pigments and other compounds. Recently, however, non-conventional extraction methods, which have been regarded as green extraction techniques, are the alternatives to conventional extraction since they require less amount of solvent as well as shorter extraction time and are more environment-friendly (Cheok et al., 2014). Selecting an appropriate extraction technique for each type of natural pigment must be done to improve the efficiency and productivity of natural colorant. The effects of selected extraction methods on the stability and extraction yield of pigments are discussed here to serve as a guideline for the proper selection of extraction technique.

Conventional extraction methods

Conventional methods, such as Soxhlet extraction, maceration, and hydrodistillation, are simple, inexpensive and easy to handle (Veggi et al., 2013). These methods have, therefore, been widely used to extract essential oils, bioactive compounds, and natural pigments from a wide array of plant materials. Soxhlet extraction was initially designed for lipid extraction but now it is also used as a reference method for comparing the yield of an advanced extraction technique (Azmir et al., 2013). During the

extraction, a plant sample contained in a thimble is repeatedly percolated with condensed vapor of solvent until the extraction is completed (or the solvent is no longer able to solubilize interested compounds in the sample), which is noted when the solvent has become colorless (Veggi et al., 2013). Although Soxhlet extraction is simple and easy to handle, it requires large amount of solvent, long extraction time, and leads to pigment degradation due to heat.

Maceration is a conventional extraction method that can be performed at room temperature. This method can thus be used for extracting heat-sensitive pigments. Cha et al. (2010a), for instance, noted that the pheophytin content, which is an indirect indicator of chlorophyll degradation, of a crude extract of green microalgae (*Chlorella vulgaris*) obtained from maceration at 25°C for 6 h was lower than that from Soxhlet extraction at 100°C for 2 h. To extract crude pigments, a plant sample is ground and mixed with an extraction solvent, and the mixture is left in an extraction vessel with occasional shaking or stirring. After the process is finished, the liquid is strained off and the residue is pressed by a mechanical press, or centrifuged to repeat extraction with a fresh solvent until the solvent exhibits no color (Azmir et al., 2013). Since the extraction is generally performed at room temperature, the time required for the extraction is long; large amount of solvent is also needed to repeat the extraction.

Hydrodistillation is a traditional method that is usually used for extracting essential oils from a plant material. The extraction medium, which is hot water and/or steam, is

directly in contact of a sample to extract interested compounds (Azmir et al., 2013). Since hydrodistillation involves the use of water for extraction, an obtained extract is not contaminated with any organic solvent, which can be toxic. However, since hydrodistillation is normally operated at a rather higher temperature (about 100°C), significant pigment degradation is expected.

The efficiency of conventional extraction methods directly depends on the solubility of a solute from a plant material into an extraction solvent (Cowan, 1999); extraction temperature also plays an important role on extraction efficiency. Examples of solvents used for extracting chlorophylls, carotenoids, betalains, and anthocyanins are listed in Table 3. Although conventional extraction methods have many advantages, these require large amount of solvent, long extraction time, and may lead to significant pigment degradation (Cheok et al., 2014). Non-conventional methods or green extraction methods have therefore been proposed to alleviate such limitations.

Non-conventional extraction methods

Supercritical fluid extraction

Supercritical fluid extraction utilizes the advantages of supercritical fluids, which exhibit gas- and liquid-like properties to enhance extraction. A supercritical fluid can be produced by subjecting a solvent to a temperature and pressure beyond its critical point. In supercritical state, fluid possesses high diffusivity and low viscosity similar to gas but exhibits high solvation power similar to liquid (Macías-Sánchez et al., 2005). For these reasons, supercritical fluids can better penetrate into a sample matrix; this subsequently leads to a more efficient extraction.

Carbon dioxide is considered an ideal solvent for SFE since its critical temperature (T_c) and critical pressure (P_c), 31°C and 74 bars, respectively, are not too high (Greibrokk, 1991). SFE is generally operated within a pressure range of 8 to 40 MPa and a temperature range of 30 to 60°C. For this reason, SFE can be applied to extract heat-sensitive pigments. Vági et al. (2002) reported that the pheophytin contents of crude extracts of *Origanum majorana* L. obtained from SFE at 40, 50, and 60°C were lower than those obtained from Soxhlet extraction at 70 and 80°C.

Since most extraction solvents for SFE are non-polar, this method is suitable for the extraction of low-polar pigments such as carotenoids and chlorophylls. On the other hand, SFE is not suitable for extraction of betalains and anthocyanins, which are high-polar pigments. Macías-Sánchez et al. (2005) reported that SFE was faster and more selective than ultrasound-assisted maceration for extracting carotenoids and

chlorophylls from *Nannochloropsis gaditana*. The highest extraction yields of both carotenoids and chlorophyll *a* via SFE were achieved when using an extraction temperature of 60°C and a pressure of 400 bar. Nevertheless, SFE with supercritical carbon dioxide as a solvent resulted in lower carotenoids and chlorophyll *a* yields than ultrasound-assisted maceration employing methanol as a solvent.

To improve the extraction yields of carotenoids and chlorophylls, a combination of carbon dioxide and organic solvent (e.g., methanol, ethanol) has been proposed. Addition of 5% (v/v) ethanol to supercritical carbon dioxide was noted to increase the extraction yield of β -carotene in carrot by about 7% (Baysal et al., 2000). Using 7% (v/v) ethanol as a co-solvent with supercritical carbon dioxide could also increase the extraction yield of chlorophylls *a* and *b* from 0.027 and 0.023 to 0.848 and 0.356 mg/g dry sample, respectively (Guedes et al., 2013).

Although SFE is quite efficient and involves small solvent consumption, no or less use of toxic solvent can extract heat-sensitive pigments and can be automated. This extraction method requires high capital and operating costs because of the high pressure required for operation (Veggi et al., 2013).

Pressurized liquid extraction

Pressurized liquid extraction utilizes a liquid solvent at elevated pressure (10.3–13.8 MPa) and temperature (40–200°C) for extraction (Antunes et al., 2008). High temperature results in the better diffusion of solvent into sample matrix and also helps disrupt plant cells, resulting in a more effective release of pigments from the cells, and hence more effective extraction. High pressure, on the other hand, forces solvent into matrix pores and hence allows better contact between the solvent and compounds to be extracted (Cha et al., 2010b; Mustafa and Turner, 2011). Thus, PLE requires shorter time and involves the use of less solvent for extraction (Antunes et al., 2008).

Pressurized liquid extraction can extract both water- and oil-based pigments, depending on the selection of an extraction solvent. However, PLE cannot effectively extract heat-sensitive pigments since the method involves the use of high temperature. PLE is therefore normally used to extract less heat-sensitive pigments (i.e., anthocyanins, carotenoids, and chlorophylls). Machado et al. (2015) studied the use of PLE in comparison with conventional methods (Soxhlet extraction and maceration) to extract monomeric anthocyanins from blackberry. PLE at 100°C exhibited a higher extraction rate than Soxhlet extraction at 80°C and maceration extraction at 25°C. Time required to extract monomeric anthocyanins in the case of PLE, Soxhlet extraction, and maceration was 30, 300, and 1440 min, respectively. In terms of carotenoids and chlorophylls, Cha et al. (2010a) reported that PLE at 160°C for 120 min yielded higher contents of carotenoids and chlorophylls *a* and *b* than maceration extraction at 25°C for 360 min and Soxhlet extraction at 100°C for 120 min. Although PLE gave the highest yields of carotenoids and chlorophylls *a* and *b*, the method resulted in the highest pheophytin content due to the use of higher temperature for extraction.

To increase the stability of pigments during PLE, some pretreatment methods, which are mentioned in the section “Stability Enhancement via Pretreatments,” can be used. Adjusting the pH of an extraction solvent to acidic values can help retard

Table 3. Examples of solvents used for extracting different groups of pigments (Cowan, 1999; Kujala et al., 2001; Hosikian et al., 2010; Machmudah and Goto, 2013).

Chlorophylls	Carotenoids	Betalains	Anthocyanins
Methanol	Methanol	Methanol	Methanol
Ethanol	Ethanol	Water	Water
Acetone	Acetone		
	n-hexane		
	Pentane		
	Chloroform		

the degradation of anthocyanins during PLE because acidic condition could lead to the formation of flavylium cation, which is the most stable form of anthocyanin (Tan et al., 2014). Sharif et al. (2010) reported that maximum redness (or the highest a^* value) and the highest yield of anthocyanins (cyanidin 3-glucoside, cyaniding 3-rutinoside, and cyanidin chloride) extracted from onion (*Allium cepa*) skin could be achieved by adding 0.1% (v/v) hydrochloric acid into an extraction solvent (methanol) and performing PLE at a temperature of 80°C and pressure of 689.48 bar. Besides adjusting the pH, intra- and intermolecular co-pigmentation may be used to increase the stability of anthocyanins during PLE. For chlorophylls, the conversion of native chlorophylls to metallo-chlorophyll derivatives prior to pigment extraction maybe employed to retain the green color of chlorophyll extracts obtained by PLE. Nevertheless, this extraction method again requires high capital and operating costs due to the use of higher pressure for extraction.

Microwave-assisted extraction

Microwave-assisted extraction is gaining interest as an alternative for extracting both water- and oil-based plant pigments due to its speed and small solvent consumption when compared with conventional extraction methods (Dahmoune et al., 2014). Rapid heating by microwave radiation results in an expansion of plant cell structure with subsequent rupture of plant cell walls. Thus, compounds, including pigments, can easily migrate out of cells, resulting in an enhanced extraction rate (Zou et al., 2013). Dabiri et al. (2005), for example, illustrated that MAE could reduce the time and solvent volume requirement when extracting pigments (alizarin and purpurin) from Rubiaceae plants. MAE required the time and solvent volume of only 20 min and 20 mL, respectively, while Soxhlet extraction required the time and solvent volume of up to 360 min and 100 mL, respectively. Moreover, MAE at an optimized condition yielded higher alizarin and purpurin recovery than Soxhlet extraction operated at its optimum condition.

Combination of stability enhancement and pigment extraction has been proposed for MAE. Cardoso-Ugarte et al. (2014) proposed a two-stage MAE with the addition of L-ascorbic acid into 50% (v/v) ethanol for the extraction of betanines and betaxanthins from red beet. Both extraction steps were performed at a power level of 400 W and 100% duty cycles. A cooling step was added between the first and second stages to delay the degradation of the pigments already extracted. The addition of ascorbic acid could retard the degradation of betanines. However, the stability of betaxanthins could not be retained by the addition of ascorbic acid, since betaxanthins are not stable in acidic condition (Stintzing and Carle, 2008).

Combination of MAE with vacuum, resulting in the so-called vacuum microwave-assisted extraction (VMAE), has recently been proposed to extract heat-sensitive bioactive compounds and pigments (Hiranvarachat et al., 2015). VMAE can reduce pigment degradation by oxidation since less oxygen is available in the extraction process. Xiao et al. (2012) noted that VMAE could prevent the degradation of vitamin C extracted from peppers and guava, β -carotene extracted from carrot, and aloin A extracted from aloe vera. The extraction yields of vitamin C, β -carotene, and aloin A extracted by VMAE were also noted to be higher than those obtained via MAE.

Ultrasound-assisted extraction

Ultrasound-assisted extraction utilizes cavitation bubbles created by ultrasound waves to enhance extraction efficiency (Rastogi, 2011). Cavitation bubbles are formed when ultrasound waves pass through a medium, creating alternative compression and decompression cycles, which in turn result in the compression and expansion of bubbles. When bubbles grow too large to be contained by the surface tension force, they collapse, resulting in shearing forces to break up or disrupt cell walls of a contacted plant material (Pitt et al., 2004). As a result, release of intracellular compounds is enhanced. UAE has received considerable attention because of its benefits, which include the ability to perform extraction at lower temperatures due to the absence of external input heat and at low solvent consumption (Tao and Sun, 2015).

Both water- and oil-based pigments can be extracted by UAE. Since UAE is a non-thermal process, it can be used to extract heat-sensitive pigments. Tao et al. (2014), for example, investigated the use of UAE to extract anthocyanins from wine lees in comparison with the use of maceration extraction, and found that at the same extraction time and temperature (36 min and 60°C), the yield of anthocyanins extracted by ultrasound at 40 kHz was higher than that by maceration. UAE has also been employed as an efficient technique for the extraction of betalains and chlorophylls. Kong et al. (2012), for example, reported that UAE could increase extraction rate by up to 88.9% in comparison with that of maceration when extracting chlorophylls from *Chlorella vulgaris*.

Ultrasound-assisted extraction is known for its lower extraction effectiveness when compared with some other non-conventional extraction processes (i.e., SFE, PLE, MAE, and PEF extraction). Cha et al. (2010a), for instance, compared the extraction yields of carotenoids and chlorophylls from *Chlorella vulgaris* by PLE and UAE and noted that PLE at 160°C for 30 min gave higher yields of both carotenoids and chlorophylls than UAE at 25°C for 360 min. Nevertheless, because of its higher temperature operation, PLE caused more chlorophyll degradation, which resulted in higher pheophytin content in the extract. To increase the extraction yield of UAE, repeated extraction is required; larger volume of solvent would be needed to repeat the extraction, however.

Recently, combined use of UAE and MAE is proposed to increase the extraction yield of UAE and, at the same time, reduce the time a material needs to undergo MAE. UAE and MAE in combination lead to more extensive damage to plant structure than employing UAE or MAE alone (Pongmalai et al., 2015). Lianfu and Zelong (2008) reported that UAE+MAE at 98 W of microwave power with 40 kHz of ultrasound gave higher yield of extracted lycopene from tomato than UAE alone. The proportion of lycopene yield was 97.4% and 89.4% for UAE+MAE and UAE, respectively. Moreover, UAE+MAE required only six minutes to obtain such a yield, while UAE needed as long as 29 min to obtain a similar yield.

Pulsed-electric field (PEF) extraction

Pulsed electric field (PEF) has been noted to be useful for enhancing many processes of food production, including the extraction process. Short- and high-voltage electric field is used to induce pore formation in the cell walls of a plant material, which

Table 4. Natural pigments extracted via different methods.

Extraction technique	Plant material	Operating conditions	Extracted pigment and yield (in parentheses) ($\mu\text{g}/\text{mg}$ sample)	Reference
SFE	Carrot (<i>Daucus carota</i> L.)	Temperature: 40°C, Pressure: 505 bar, Solvent: CO ₂ , Flow rate: 600–750 mL/min, Time: 60 min	α -carotene (0.12) β -carotene (0.15)	Chandra and Nair, 1997
SFE	<i>Dunaliella salina</i>	Temperature: 60°C, Pressure: 300 bar, Solvent: CO ₂ , Flow rate: 4.5 mmol/min, Time: 180 min	Carotenoids (14.92) chlorophylls (0.27)	Macías-Sánchez et al., 2009
SFE	<i>Scenedesmus obliquus</i>	Temperature: 40°C, Pressure: 250 bar, Solvent: CO ₂ with 7.7% (v/v) ethanol, Flow rate: 2 g/min, Time: 240 min	Carotenoids (0.30), Chlorophyll <i>a</i> (0.85), Chlorophyll <i>b</i> (0.36), Chlorophyll <i>c</i> (0.02)	Guedes et al., 2013
PLE	<i>Chlorella vulgaris</i>	Temperature: 160°C, Pressure: 103.42 bar, Solvent: 90% ethanol, Time: 30 min	β -carotene (0.50), Chlorophyll <i>a</i> (9.63), Chlorophyll <i>b</i> (5.77), Pheophytin <i>a</i> (5.64)	Cha et al., 2010a
UAE	<i>Chlorella vulgaris</i>	Temperature: room temp., Solvent: 90% ethanol, Solvent volume: 50 mL, Time: 120 min	β -carotene (0.10), Chlorophyll <i>a</i> (5.12), Chlorophyll <i>b</i> (3.17), Pheophytin <i>a</i> (2.64)	
Soxhlet extraction	<i>Chlorella vulgaris</i>	Temperature: 100°C, Solvent: 90% ethanol, Solvent volume: 100 mL, Time: 120 min	β -carotene (0.26), Chlorophyll <i>a</i> (3.32), Chlorophyll <i>b</i> (3.45), Pheophytin <i>a</i> (3.90)	
Maceration	<i>Chlorella vulgaris</i>	Temperature: room temp., Solvent: 90% ethanol, Solvent volume: 50 mL, Time: 360 min	β -carotene (0.08), Chlorophyll <i>a</i> (4.26), Chlorophyll <i>b</i> (2.58), Pheophytin <i>a</i> (2.31)	
PLE	Blackberry (<i>Rubus fruticosus</i> L.)	Temperature: 100°C, Pressure: 75 bar, Solvent: water with 50% (v/v) ethanol, Flow rate: 3.35 mL/min, Time: 30 min	Anthocyanins (1.02)	Machado et al., 2015
Soxhlet extraction	Blackberry (<i>Rubus fruticosus</i> L.)	Temperature: 80°C, Solvent: Methanol, Solvent volume: 200 mL, Time: 300 min	Anthocyanin (1.33)	
Soxhlet extraction	Blackberry (<i>Rubus fruticosus</i> L.)	Temperature: 80°C, Solvent: Ethanol, Solvent volume: 200 mL, Time: 300 min	Anthocyanin (1.68)	
Maceration	Blackberry (<i>Rubus fruticosus</i> L.)	Temperature: room temp. Solvent: methanol with 0.01% (v/v) HCl, Solvent volume: 200 mL, Time: 1440 min	Anthocyanin (1.21)	
MAE	Purple corn (<i>Zea mays</i> L.)	Solvent: 95% ethanol with 1.5-M HCl, Solvent to sample ratio (mL/g): 20:1, Microwave power: 555 W, Time: 19 min	Anthocyanins (0.18)	Yang and Zhai, 2010
Maceration	Purple corn (<i>Zea mays</i> L.)	Solvent: 95% ethanol with 1.5-M HCl, Solvent to sample ratio (mL/g): 30:1, Temperature: 55°C, Time: 60 min	Anthocyanins (0.16)	
MAE	Red beet (<i>Beta vulgaris</i>)	Solvent: 50% (v/v) ethanol with 0.04-M ascorbic acid, Solvent to sample ratio (mL/g): 250:1, Microwave power: 400 W, Time: 3.50 min	Betanines (0.19)	Cardoso-Ugarte et al., 2014
Maceration	Red beet (<i>Beta vulgaris</i>)	Solvent: 50% (v/v) ethanol, Solvent to sample ratio (mL/g): 250:1, Temperature: 80°C, Time: 60 min	Betanines (0.15)	
MAE	Carrot	Temperature: 58°C, Solvent: mixed solvent (50% v/v hexane, 25% v/v acetone, 25% v/v ethanol), Solvent volume: 75 mL, Microwave power: 180 W, Time: 3 min	Total carotenoids (0.52), β -carotene (0.23)	Hiranvarachat et al., 2013
Soxhlet extraction	Carrot	Temperature: 58°C, Solvent: mixed solvent (50% v/v hexane, 25% v/v acetone, 25% v/v ethanol), Solvent volume: 75 mL, Time: 360 min	Total carotenoids (0.61), β -carotene (0.29)	
UAE (Ultrasonic probe)	<i>Bougainvillea glabra</i> flower	Temperature: 55°C, Solvent: 50% methanol, Solvent to sample ratio (mL/g): 17:1, Power and frequency: 88 W, 20 kHz, Time: 37 min	Betacyanin (1.72) Betaxanthin (5.78)	Maran et al., 2015
UAE (Ultrasonic bath)	<i>Chlorella vulgaris</i>	Temperature: 50°C, Solvent: 80% ethanol, Solvent to sample ratio (mL/g): 200:1, Power and frequency: 200 W, 40 kHz, Time: 30 min	Total chlorophylls (13.2)	Kong et al., 2012
Maceration	<i>Chlorella vulgaris</i>	Temperature: 50°C, Solvent: 80% ethanol, Solvent to sample ratio (mL/g): 200:1, Time: 30 min	Total chlorophylls (8.32)	
UAE (Ultrasonic bath)	Light wine lees (60% Cabernet Sauvignon, 30% Merlot, 10% Cabernet Franc)	Temperature: 60°C, Solvent: 50% ethanol, Solvent to sample ratio (mL/g): 60:1, Acoustic energy density: 48 W/L, Frequency: 40 kHz, Time: 36 min	Total anthocyanins (6.69)	Tao et al., 2014
Maceration	Light wine lees (60% Cabernet Sauvignon, 30% Merlot, 10% Cabernet Franc)	Temperature: 60°C, Solvent: 50% ethanol, Solvent to sample ratio (mL/g): 60:1, Time: 36 min	Total anthocyanins (5.55)	
PEF-assisted extraction	Grape by-product	PEF treatment Temperature: room temp., Number of pulses: 30, Pulse frequency: 2 Hz, Electric field strength: 9 kV/cm, Extraction (maceration) temperature: 70°C, Solvent: 50% (v/v) ethanol, Solvent to sample ratio (mL/g): 4.5:1, Time: 60 min	Anthocyanins (14.05)	Corrales et al., 2008

(Continued on next page)

Table 4. (Continued)

Extraction technique	Plant material	Operating conditions	Extracted pigment and yield (in parentheses) ($\mu\text{g}/\text{mg}$ sample)	Reference
UAE (Ultrasonic bath)	Grape by-product	Temperature: 70°C, Solvent: 50% (v/v) ethanol, Solvent to sample ratio (mL/g): 4.5:1, Frequency: 35 kHz, Time: 60 min	Anthocyanins (7.76)	
Maceration	Grape by-product	Temperature: 70°C, Solvent: 50% (v/v) ethanol, Solvent to sample ratio (mL/g): 20:1, Time: 180 min	Anthocyanins (7.93)	Corrales et al., 2008
PEF-assisted extraction	Red beetroot	PEF treatment Temperature: 30°C, Number of pulses: 5, Pulse frequency: 1 Hz, Electric field strength: 7 kV/cm, Extraction (maceration) temperature: 30°C, Solvent: McIlvaine buffer at pH 3.5, Solvent volume: 400 mL, Time: 300 min	Betanines (90% of total)	López et al., 2009
Maceration	Red beetroot	Temperature: 50°C, Solvent: McIlvaine buffer at pH 3.5, Solvent volume: 400 mL, Time: 400 min	Betanines (66% of total)	
EAE	Marigold flower (<i>Tagetes erecta</i>)	Enzyme treatment, Enzyme to sample ratio (mL/g): 0.1:1, Time: 60 min, Extraction (maceration) temperature: 25°C, Solvent: Hexane, Solvent to sample ratio (mL/g): 4:1, Time: 1440 min	Carotenoids (0.5)	Barzana et al., 2002

subsequently leads to better release of cellular constituents, and hence enhanced extraction (Donsì et al., 2010; Azmir et al., 2013).

Most solvents used to extract betalains and anthocyanins are polar solvents, possessing electrical conductivity, and can let electricity pass thorough to sample cells. On the other hand, electric field cannot pass thorough a non-polar solvent, since it is an electrical resistance possessing low or negligible conductivity (Yuhas, 1995). For this reason, PEF is more suitable for the extraction of betalains and anthocyanins than for chlorophylls and carotenoids.

A number of works exist on the use of PEF to extract (or to assist the extraction of) plant pigments. Puértolas et al. (2013), for instance, applied PEF at 3.4 kV/cm and 105 μs (35 pulses of 3 μs) to extract anthocyanins from purple-fleshed potato. The anthocyanins yield was noted, as expected, to be higher than those from the non-PEF-treated sample and also from the macerated sample. pH value also exerts a significant effect on the PEF extraction yield. López et al. (2009), for example, studied the effect of pH of the solvent on the PEF extraction yield of betanines from red beetroot, and found that McIlvaine buffer at pH 3.5 led to the highest yield of betanines. This is probably because an acidic solvent could prevent the degradation of betanines during extraction (Bilyk and Howard, 1982).

Notwithstanding its potential, PEF-based extractor, especially at an industrial scale, suffers from the need of a high-power supply equipment and treatment chamber (Nip, 2007), which are nowadays still rather expensive.

Enzyme-assisted extraction

Enzyme-assisted extraction involves the use of enzymes to enhance the extraction of bioactive compounds, including pigments, from a plant material. Various enzymes, such as pectinase, cellulase, and hemicellulase, are used to hydrolyze plant cellulosic cell walls (Socaciu, 2008), resulting in easier release of cellular constituents, and hence facilitated extraction. EAE is of interest due to its non-thermal nature as well as lower toxicity and solvent consumption (Socaciu, 2008). EAE can be divided into two common approaches, which are enzyme-assisted

aqueous extraction (EAAE) and enzyme-assisted cold pressing (EACP) (Latif and Anwar, 2009). In EAAE, enzymes are used to help destroy cell walls, and hence rupturing polysaccharide-protein colloids. On the other hand, enzymes are used only to hydrolyze cell walls in the case of EACP (Hassan and Gökçe, 2014).

Many researchers investigated the effect of enzyme treatment on the extraction yields of chlorophylls, carotenoids, betalains, and anthocyanins. Barzana et al. (2002), for example, noted that the extraction yield of carotenoids from marigold flower treated by enzymes (viscozyme, proteolytic, and pectinolytic enzymes) was higher than that from the untreated sample. Kammerer et al. (2005) proposed a two-stage process for the extraction of anthocyanins from grape pomace via EAE. The process involved a pre-extraction step, which was then followed by EAE. Release of phenolic acids during the pre-extraction step nevertheless caused a decrease in pH, leading to an unfavorable condition for enzyme activity. For this reason, before EAE, the pH of the residue obtained in the pre-extraction step was adjusted to an optimum range for enzyme hydrolysis, which resulted in enhanced extraction.

Enzyme treatment can be also applied to assist other green extraction methods. Lenucci et al. (2015), for instance, proposed an enzyme-aided supercritical carbon dioxide extraction, and found that enzymatic pretreatment could help increase the extractable lycopene concentration from the matrix of tomato puree due to the degradation of tomato cell walls by enzymes. However, the denseness of the enzyme-treated sample hindered the diffusion of supercritical carbon dioxide through the sample matrix. Addition of a co-matrix (hazelnut seeds) into the base matrix was therefore proposed to facilitate the penetration of supercritical carbon dioxide through the sample matrix, and hence increased lycopene extractability. Although EAE can improve extraction yield and can be applied to extract heat-sensitive pigments, EAE requires a very long extraction time; enzymes are also normally quite expensive (Hardouin et al., 2014).

Examples of pigment extraction via different methods, and pros and cons of each extraction method are listed in Tables 4 and 5, respectively.

Table 5. Advantages and drawbacks of different extraction methods.

Extraction method	Solvent	Advantages	Drawbacks
Soxhlet extraction	<ul style="list-style-type: none"> Organic solvents (both polar and non-polar) 	<ul style="list-style-type: none"> Easy to handle Low investment Automated system 	<ul style="list-style-type: none"> Not suitable for heat-sensitive pigments Long extraction time
Maceration	<ul style="list-style-type: none"> Organic solvents (both polar and non-polar) 	<ul style="list-style-type: none"> Easy to handle Low investment Applicable for heat-sensitive pigments 	<ul style="list-style-type: none"> Very long extraction time Large solvent consumption Filtration step required Non-automated system Repeated extraction may be required Not suitable for heat-sensitive pigments Long extraction time
Hydrodistillation	<ul style="list-style-type: none"> Water 	<ul style="list-style-type: none"> Easy to handle Low investment Automated system 	<ul style="list-style-type: none"> Required sophisticated safety controls High pressure leading to high capital and operating costs
SFE	<ul style="list-style-type: none"> CO₂ (non-polar) CO₂ with polar solvent 	<ul style="list-style-type: none"> Moderate extraction time Applicable for heat-sensitive pigments No use of toxic solvents Automated system 	<ul style="list-style-type: none"> Not suitable for heat-sensitive pigments High investment
PLE	<ul style="list-style-type: none"> Organic solvents (both polar and non-polar) 	<ul style="list-style-type: none"> Fast extraction Small solvent consumption Automated system 	<ul style="list-style-type: none"> Not suitable for heat-sensitive pigments High investment
MAE	<ul style="list-style-type: none"> Organic solvents (both polar and non-polar) 	<ul style="list-style-type: none"> Fast extraction Easy to handle Moderate solvent consumption Moderate investment 	<ul style="list-style-type: none"> Not suitable for heat-sensitive pigments Filtration step required Non-automated system
PEF extraction	<ul style="list-style-type: none"> Organic solvents (both polar and non-polar) 	<ul style="list-style-type: none"> Fast extraction Applicable for heat-sensitive pigments Moderate investment 	<ul style="list-style-type: none"> Large solvent consumption Filtration step required Non-automated system
UAE	<ul style="list-style-type: none"> Organic solvents (both polar and non-polar) 	<ul style="list-style-type: none"> Fast extraction Applicable for heat-sensitive pigments Simple Low investment 	<ul style="list-style-type: none"> Large solvent consumption Filtration required Non-automated system Repeated extraction may be required
EAE	<ul style="list-style-type: none"> Organic solvents (both polar and non-polar) 	<ul style="list-style-type: none"> Applicable for heat-sensitive pigments Easy to handle Moderate solvent consumption 	<ul style="list-style-type: none"> Long extraction time Filtration required Non-automated system

Conclusions

Selected methods that can be used to enhance the stability of natural pigments during their processing and storage were first outlined. Selection of an appropriate pretreatment method depends on the type of pigments. Selected non-conventional extraction methods are then reviewed as a means to enhance yield and maintain the stability of pigments without sacrificing friendliness to the environment. Suitability of extraction methods is discussed based on the thermal sensitivity and polarity of pigments. MAE seems to be a proper method for the extraction of carotenoids, chlorophylls, and anthocyanins but not heat-sensitive pigments such as betalains. UAE, EAE, and PEF, which are classified as non-thermal extraction processes, can be used to extract betalains.

Although only guidelines for the selection of appropriate pretreatment and extraction methods for each type of pigment are given, other factors such as investment cost, operating cost as well as applicability of the selected techniques at an industrial scale must also be considered besides extraction yield and stability of pigments. Combination of different extraction and stability enhancement methods can be performed to increase both pigment stability and extraction yield.

Funding

The authors express their sincere appreciation to the Thailand Research Fund (TRF, Grant No. RTA 5880009) for supporting the study financially.

ORCID

Sakamon Devahastin  <http://orcid.org/0000-0001-9582-1554>

References

- Ahmed, J., Al-Salmam, F. and Almusallam, A. S. (2013). Effect of blanching on thermal color degradation kinetics and rheological behavior of rocket (*Eruca sativa*) puree. *J. Food Eng.* **119**:660–667.
- Ananga, A., Georgiev, V., Ochieng, J., Phills, B. and Tsolova, V. (2013). Production of anthocyanins in grape cell cultures: A potential source of raw material for pharmaceutical, food, and cosmetic industries. In: *The Mediterranean Genetic Code – Grapevine and Olive*, pp. 247–287. Poljuha, D. and Sladonja, B., Eds., InTech, Rijeka.
- Antunes, P., Viana, P., Vinhas, T., Capelo, J. L., Rivera, J. and Gaspar, E. M. S. M. (2008). Optimization of pressurized liquid extraction (PLE) of dioxin-furans and dioxin-like PCBs from environmental samples. *Talanta*. **75**:916–925.
- Azmir, J., Zaidul, I. S. M., Rahman, M. M., Sharif, K. M., Mohamed, A., Sahena, F., Jahuul, M. H. A., Ghafoor, K., Norulaini, N. A. N. and Omar, A. K. M. (2013). Techniques for extraction of bioactive compounds from plant materials: A review. *J. Food Eng.* **117**:426–436.
- Bahçeci, K. S., Serpen, A., Gökmen, V. and Acar, J. (2005). Study of lipoxigenase and peroxidase as indicator enzymes in green beans: Change of enzyme activity, ascorbic acid and chlorophylls during frozen storage. *J. Food Eng.* **66**:187–192.
- Bąkowska-Barczak, A. (2005). Acylated anthocyanins as stable, natural food colorants – A review. *Pol. J. Food Nutr. Sci.* **14**:107–116.
- Bąkowska, A., Kucharska, A. Z. and Oszmiański, J. (2003). The effects of heating, UV irradiation, and storage on stability of the anthocyanin – polyphenol copigment complex. *Food Chem.* **81**:349–355.
- Barzana, E., Rubio, D., Santamaria, R. I., Garcia-Correa, O., Garcia, F., Ridaura Sanz, V. E. and López-Munguía, A. (2002). Enzyme-mediated solvent extraction of carotenoids from marigold flower (*Tagetes erecta*). *J. Agric. Food Chem.* **50**:4491–4496.
- Baysal, T., Ersus, S. and Starmans, D. A. J. (2000). Supercritical CO₂ extraction of β -carotene and lycopene from tomato paste waste. *J. Agric. Food Chem.* **48**:5507–5511.
- Bilyk, A. and Howard, M. (1982). Reversibility of thermal degradation of betacyanines under the influence of isoascorbic acid. *J. Agric. Food Chem.* **30**:906–908.

- Bilyk, A., Kolodij, M. A. and Sapers, G. M. (1981). Stabilization of red beet pigments with isoascorbic acid. *J. Food Sci.* **46**:1616–1617.
- Bobbio, P. A. and Guedes, M. C. (1990). Stability of copper and magnesium chlorophylls. *Food Chem.* **36**:165–168.
- Boon, C. S., McClements, D. J., Weiss, J. and Decker, E. A. (2010). Factors influencing the chemical stability of carotenoids in foods. *Crit. Rev. Food Sci. Nutr.* **50**:515–532.
- Bordignon-Luiz, M. T., Gauche, C., Gris, E. F. and Falcão, L. D. (2007). Colour stability of anthocyanins from Isabel grapes (*Vitis labrusca* L.) in model systems. *LWT Food Sci. Technol.* **40**:594–599.
- Britton, G. (1996). Carotenoids. In: *Natural Food Colorants*, 2nd ed., pp. 197–243. Hendry, G. A. F. and Houghton, J. D., Eds., Chapman & Hall, New York, NY.
- Buchweitz, M., Brauch, J., Carle, R. and Kammerer, D. R. (2013). Application of ferric anthocyanin chelates as natural blue food colorants in polysaccharide and gelatin based gels. *Food Res. Int.* **51**:274–282.
- Buchweitz, M., Nagel, A., Carle, R. and Kammerer, D. R. (2012). Characterisation of sugar beet pectin fractions providing enhanced stability of anthocyanin-based natural blue food colourants. *Food Chem.* **132**:1971–1979.
- Cardoso-Ugarte, G. A., Sosa-Morales, M. E., Ballard, T., Liceaga, A. and Martín-González, M. F. S. (2014). Microwave-assisted extraction of betalains from red beet (*Beta vulgaris*). *LWT Food Sci. Technol.* **59**:276–282.
- Castañeda-Ovando, A., Pacheco-Hernández, M. D. L., Páez-Hernández, M. E., Rodríguez, J. A. and Galán-Vidal, C. A. (2008). Chemical studies of anthocyanins: A review. *Food Chem.* **113**:859–871.
- Cevallos-Casals, B. A. and Cisneros-Zevallos, L. (2004). Stability of anthocyanin-based aqueous extracts of Andean purple corn and red-fleshed sweet potato compared to synthetic and natural colorants. *Food Chem.* **86**:69–77.
- Cha, K. H., Lee, H. J., Koo, S. Y., Song, D. G., Lee, D. U. and Pan, C. H. (2010a). Optimization of pressurized liquid extraction of carotenoids and chlorophylls from *Chlorella vulgaris*. *J. Agric. Food Chem.* **58**:793–797.
- Cha, K. H., Kang, S. W., Kim, C. Y., Um, B. H., Na, Y. R. and Pan, C. H. (2010b). Effect of pressurized liquids on extraction of antioxidants from *Chlorella vulgaris*. *J. Agric. Food Chem.* **58**:4756–4761.
- Chandra, A. and Nair, M. G. (1997). Supercritical fluid carbon dioxide extraction of α - and β -carotene from carrot (*Daucus carota* L.). *Phytochem. Anal.* **8**:244–246.
- Chandrasekhar, J., Sonika, G., Madhusudhan, M. C. and Raghavarao, K. S. M. S. (2015). Differential partitioning of betacyanins and betaxanthins employing aqueous two-phase extraction. *J. Food Eng.* **144**:156–163.
- Cheok, C. Y., Salman, H. A. K. and Sulaiman, R. (2014). Extraction and quantification of saponins: A review. *Food Res. Int.* **59**:16–40.
- Choi, M. H., Kim, G. H. and Lee, H. S. (2002). Effects of ascorbic acid retention on juice color and pigment stability in blood orange (*Citrus sinensis*) juice during refrigerated storage. *Food Res. Int.* **35**:753–759.
- Corrales, M., Toepfl, S., Butz, P., Knorr, D. and Tauscher, B. (2008). Extraction of anthocyanins from grape by-products assisted by ultrasonics, high hydrostatic pressure or pulsed electric fields: A comparison. *Innovative Food Sci. Emerging Technol.* **9**:85–91.
- Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clin. Microbiol. Rev.* **12**:564–582.
- Dabiri, M., Salimi, S., Ghassempour, A., Rassouli, A. and Talebi, M. (2005). Optimization of microwave-assisted extraction for alizarin and purpurin in Rubiaceae plants and its comparison with conventional extraction methods. *J. Sep. Sci.* **28**:387–396.
- Dahmoune, F., Moussia, S. K., Reminia, H., Cherbac, A. and Madania, K. (2014). *Pistacia lentiscus* leaves as a source of phenolic compounds: Microwave-assisted extraction optimized and compared with ultrasound-assisted and conventional solvent extraction. *Ind. Crops Prod.* **61**:31–40.
- Delgado-Vargas, F., Jiménez, A. R. and Paredes-López, O. (2000). Natural pigments: Carotenoids, anthocyanins, and betalains – Characteristics, biosynthesis, processing, and stability. *Crit. Rev. Food Sci. Nutr.* **40**:173–289.
- Delgado-Vargas, F. and Paredes-López, O. (2003). *Natural Colorants for Food and Nutraceutical Uses*. CRC Press, Boca Raton, FL.
- Donsi, F., Ferrari, G. and Pataro, G. (2010). Application of pulsed electric fields treatments for the enhancement of mass transfer from vegetable tissue. *Food Eng. Rev.* **2**:109–130.
- Fellows, P. (2000). *Food Processing Technology – Principles and Practice*, 2nd ed. CRC Press, Boca Raton, FL.
- Francis, F. J. (1996). Safety of food colorants. In: *Natural Food Colorants*, 2nd ed., pp. 112–126. Hendry, G. A. F. and Houghton, J. D., Eds., Chapman & Hall, New York, NY.
- Gauche, C., Malagoli, E. D. S. and Bordignon-Luiz, M. T. (2010). Effect of pH on the copigmentation of anthocyanins from Cabernet Sauvignon grape extracts with organic acids. *Sci. Agric.* **67**:41–46.
- Gliemmo, M. F., Latorrea, M. E., Gerschenson, L. N. and Campos, C. A. (2009). Color stability of pumpkin (*Cucurbita moschata*, Duchesne ex Poiret) puree during storage at room temperature: Effect of pH, potassium sorbate, ascorbic acid and packaging material. *LWT Food Sci. Technol.* **42**:196–201.
- Gómez-Míguez, M., González-Manzano, S., Escribano-Bailón, M. T., Heredia, F. J. and Santos-Buelga, C. (2006). Influence of different phenolic copigments on the color of malvidin 3-glucoside. *J. Agric. Food Chem.* **54**:5422–5429.
- Gonçaves, L. C. P., Genova, B. M. D., DÖrr, F. A., Pinto, E. and Bastos, E. L. (2013). Effect of dielectric microwave heating on the color and anti-radical capacity of betanine. *J. Food Eng.* **118**:49–55.
- Greibrokk, T. (1991). Fluid extraction of chlorinated compounds and other pollutants. In: *Organic Micropollutants in the Aquatic Environment*, pp. 112–114. Angeletti, G. and Bjørseth, A., Eds., Springer, Brussels, Belgium.
- Gris, E. F., Ferreira, E. A., Falcão, L. D. and Bordignon-Luiz, M. T. (2007). Caffeic acid copigmentation of anthocyanins from Cabernet Sauvignon grape extracts in model systems. *Food Chem.* **100**:1289–1296.
- Guedes, A. C., Gião, M. S., Matias, A. A., Nunes, A. V. M., Pintado, M. E., Duarte, C. M. M. and Malcata, F. X. (2013). Supercritical fluid extraction of carotenoids and chlorophylls a, b and c from a wild strain of *Scenedesmus obliquus* for use in food processing. *J. Food Eng.* **116**:478–482.
- Hackett, M. M., Lee, J. H., Francis, D. and Schwartz, S. J. (2004). Thermal stability and isomerization of lycopene in tomato oleoresins from different varieties. *J. Food Sci.* **69**:C536–C541.
- Han, D., Kim, S. J., Kim, S. H. and Kim, D. M. (1998). Repeated regeneration of degraded red beet juice pigments in the presence of antioxidants. *J. Food Sci.* **63**:69–72.
- Hardouin, K., Bedoux, G., Burlot, A. S., Nyvall-Collén, P. and Bourgoignon, N. (2014). Enzymatic recovery of metabolites from seaweeds: Potential applications. In: *Advances in Botanical Research*, pp. 280–313. Jacquot, J. P., Gadat, P. and Bourgoignon, N., Eds., Academic Press, Oxford, UK.
- Harp, B. P. and Barrows, J. N. (2015). US regulation of color additives in foods. In: *Colour Additives for Foods and Beverages*, pp. 75–88. Scotter, M. J., Ed., Woodhead Publishing, Cambridge.
- Hassan, E. and Gökçe, A. (2014). Production and consumption of biopesticides. In: *Advances in Plant Biopesticides*, pp. 361–380. Singh, D., Ed., Springer, New Delhi, India.
- Heaton, J. W. and Marangoni, A. G. (1996). Chlorophyll degradation in processed foods and senescent plant tissues. *Trends Food Sci. Technol.* **7**:8–15.
- Hendry, G. A. F. (1996). Natural pigments in biology. In: *Natural Food Colorants*, 2nd ed., pp. 1–39. Hendry, G. A. F. and Houghton, J. D., Eds., Chapman & Hall, New York, NY.
- Henry, B. S. (1996). Natural food colours. In: *Natural Food Colorants*, 2nd ed., pp. 40–79. Hendry, G. A. F. and Houghton, J. D., Eds., Chapman & Hall, New York, NY.
- Herbach, K. M., Rohe, M., Stintzing, F. C. and Carle, R. (2006a). Structural and chromatic stability of purple pitaya (*Hylocereus polyrhizus* [Weber] Britton & Rose) betacyanins as affected by the juice matrix and selected additives. *Food Res. Int.* **39**:667–677.
- Herbach, K. M., Stintzing, F. C. and Carle, R. (2006b). Betalain stability and degradation – Structural and chromatic aspects. *J. Food Sci.* **71**:R41–R50.
- Hiranvarachat, B., Devahastin, S. and Chiewchan, N. (2011). Effects of acid pretreatments on some physicochemical properties of carrot undergoing hot air drying. *Food Bioprod. Process.* **89**:116–127.

- Hiranvarachat, B., Devahastin, S., Chiewchan, N. and Raghavan, G. S. V. (2013). Structural modification by different pretreatment methods to enhance microwave-assisted extraction of β -carotene from carrots. *J. Food Eng.* **115**:190–197.
- Hiranvarachat, B., Devahastin, S. and Soponronnarit, S. (2015). Comparative evaluation of atmospheric and vacuum microwave-assisted extraction of bioactive compounds from fresh and dried *Centella asiatica* L. leaves. *Int. J. Food Sci. Technol.* **50**:750–757.
- Hosikian, A., Lim, S., Halim, R. and Danquah, M. K. (2010). Chlorophyll extraction from microalgae: A review on the process engineering aspects. *Int. J. Chem. Eng.* **2010**:1–11.
- Jackman, R. L. and Smith, J. L. (1996). Anthocyanins and betalains. In: *Natural Food Colorants*, 2nd ed., pp. 244–309. Hendry, G. A. F. and Houghton, J. D., Eds., Chapman & Hall, New York, NY.
- Kammerer, D., Claus, A., Schieber, A. and Carle, R. (2005). A novel process for the recovery of polyphenols from grape (*Vitis vinifera* L.) pomace. *J. Food Sci.* **70**:C157–C163.
- Khan, M. I. and Giridhar, P. (2014). Enhanced chemical stability, chromatic properties and regeneration of betalains in *Rivina humilis* L. berry juice. *LWT Food Sci. Technol.* **58**:649–657.
- Khoo, H. E., Prasad, K. N., Kong, K. W., Jiang, Y. and Ismail, A. (2011). Carotenoids and their isomers: Color pigments in fruits and vegetables. *Molecules.* **16**:1710–1738.
- Kong, W., Liu, N., Zhang, J., Yang, Q., Hua, S., Song, H. and Xia, C. (2012). Optimization of ultrasound-assisted extraction parameters of chlorophyll from *Chlorella vulgaris* residue after lipid separation using response surface methodology. *J. Food Sci. Technol.* **51**:2006–2013.
- Kujala, T., Loponen, J. and Pihlaja, K. (2001). Betalains and phenolics in red beetroot (*Beta vulgaris*) peel extracts: Extraction and characterisation. *J. Biosci.* **56**:343–348.
- Lakkakula, N., Lima, M. and Walker, T. (2004). Rice bran stabilization and rice bran oil extraction using Ohmic heating. *Bioresource Technol.* **92**:157–161.
- Latif, S. and Anwar, F. (2009). Physicochemical studies of hemp (*Cannabis sativa*) seed oil using enzyme-assisted cold-pressing. *Eur. J. Lipid Sci. Technol.* **111**:1042–1048.
- Lau, M. H., Tang, J. and Swanson, B. G. (2000). Kinetics of textural and color changes in green asparagus during thermal treatments. *J. Food Eng.* **45**:231–236.
- Lavelli, V., Zanoni, B. and Zaniboni, A. (2007). Effect of water activity on carotenoid degradation in dehydrated carrots. *Food Chem.* **104**:1705–1711.
- Lenucci, M. S., Caroli, M. D., Marrese, P. P., Iurlaro, A., Rescio, L., Böhm, V., Dalessandro, G. and Piro, G. (2015). Enzyme-aided extraction of lycopene from high-pigment tomato cultivars by supercritical carbon dioxide. *Food Chem.* **170**:193–202.
- Lianfu, Z. and Zelong, L. (2008). Optimization and comparison of ultrasound/microwave assisted extraction (UMAE) and ultrasonic-assisted extraction (UAE) of lycopene from tomatoes. *Ultrason. Sonochem.* **15**:731–737.
- López, N., Puértolas, E., Condón, S., Raso, J. and Alvarez, I. (2009). Enhancement of the extraction of betanine from red beetroot by pulsed electric fields. *J. Food Eng.* **90**:60–66.
- Machado, A. P. D. F., Pasquel-Reátegui, J. L., Barbero, G. F. and Martínez, J. (2015). Pressurized liquid extraction of bioactive compounds from blackberry (*Rubus fruticosus* L.) residues: A comparison with conventional methods. *Food Res. Int.* **77**:675–683.
- Machmudah, S. and Goto, M. (2013). Methods for extraction and analysis of carotenoids. In: *Natural Products*, pp. 3368–3407. Ramawat, K. G. and Mérillon, J. M., Eds., Springer, Heidelberg, Germany.
- Macías-Sánchez, M. D., Mantell, C., Rodríguez, M., Martínez de la Ossa, E., Lubián, L. M. and Montero, O. (2005). Supercritical fluid extraction of carotenoids and chlorophyll *a* from *Nannochloropsis gaditana*. *J. Food Eng.* **66**:245–251.
- Macías-Sánchez, M. D., Mantell, C., Rodríguez, M., Martínez de la Ossa, E., Lubián, L. M. and Montero, O. (2009). Comparison of supercritical fluid and ultrasound-assisted extraction of carotenoids and chlorophyll *a* from *Dunaliella salina*. *Talanta.* **77**:948–952.
- Maharaj, V. and Sankat, C. K. (1996). Quality changes in dehydrated dasheen leaves: Effects of blanching pre-treatments and drying conditions. *Food Res. Int.* **29**:563–568.
- Maran, J. P., Priyab, B. and Nivetha, C. V. (2015). Optimization of ultrasound-assisted extraction of natural pigments from *Bougainvillea glabra* flowers. *Ind. Crops Prod.* **63**:182–189.
- Marquez, U. M. L. and Sinnecker, P. (2008). Chlorophylls in foods: Sources and stability. In: *Food Colorants: Chemical and Functional Properties*, pp. 195–212. Socaciu, C., Ed., CRC Press, Boca Raton, FL.
- Marx, M., Stuparic, M., Schieber, A. and Carle, R. (2003). Effects of thermal processing on trans-cis-isomerization of β -carotene in carrot juices and carotene-containing preparations. *Food Chem.* **83**:609–617.
- Masone, D. and Chanforan, C. (2015). Study on the interaction of artificial and natural food colorants with human serum albumin: A computational point of view. *Comput. Biol. Chem.* **56**:152–158.
- Mazza, G. and Brouillard, R. (1990). The mechanism of co-pigmentation of anthocyanins in aqueous solutions. *Phytochemistry.* **29**:1097–1102.
- Mercadante, A. Z. and Bobbio, F. O. (2008). Anthocyanins in foods: Occurrence and physicochemical. In: *Food Colorants: Chemical and Functional Properties*, pp. 241–274. Socaciu, C., Ed., CRC Press, Boca Raton, FL.
- Mustafa, A. and Turner, C. (2011). Pressurized liquid extraction as a green approach in food and herbal plants extraction: A review. *Anal Chim. Acta.* **703**:8–18.
- Negi, P. S. and Roy, S. K. (2001). Effect of drying conditions on quality of green leaves during long-term storage. *Food Res. Int.* **34**:283–287.
- Ngo, T. and Zhao, Y. (2005). Retaining green pigments on thermally processed peels-on green pears. *J. Food Sci.* **70**:C568–C574.
- Nip, W. K. (2007). Fundamentals of food manufacturing. In: *Handbook of Food Products Manufacturing*, pp. 5–44. Hui, Y. H., Chandan, R. C., Clark, S., Cross, N., Dobbs, J., Hurst, W. J., Nollet, L. M. L., Shimoni, E., Sinha, N., Smith, E. B., Surapat, S., Titchenal, A. and Toldrá, F., Eds., Wiley, Hoboken, UK.
- Olivera, D. F., Viña, A. Z., Marani, C. M., Ferreyra, R. M., Mugridge, A., Chaves, A. R. and Mascheroni, R. H. (2008). Effect of blanching on the quality of Brussels sprouts (*Brassica oleracea* L. gemmifera DC) after frozen storage. *J. Food Eng.* **84**:148–155.
- Ötles, S. and Çagindi, Ö. (2008). Carotenoids as natural colorants. In: *Food Colorants: Chemical and Functional Properties*, pp. 51–70. Socaciu, C., Ed., CRC Press, Boca Raton, FL.
- Özkan, G. and Bilek, S. E. (2015). Enzyme-assisted extraction of stabilized chlorophyll from spinach. *Food Chem.* **176**:152–157.
- Perera, O. C. and Baldwin, E. A. (2001). Biochemistry of fruits and its implications on processing. In: *Fruit Processing: Nutrition, Products and Quality Management*, 2nd ed., pp. 25–27. Arthey, D. and Ashurst, P. R., Eds., Aspen, Gaithersburg, Germany.
- Pitt, W. G., Hussein, G. A. and Staples, B. J. (2004). Ultrasonic drug delivery – A general review. *Expert Opin. Drug Deliv.* **1**:37–56.
- Pokorny, J. (2007). Antioxidants in food preservation. In: *Handbook of Food Preservation*, 2nd ed., pp. 259–286. Rahman, M. S., Ed., CRC Press, Boca Raton, FL.
- Pongmalai, P., Devahastin, S., Chiewchan, N. and Soponronnarit, S. (2015). Enhancement of microwave-assisted extraction of bioactive compounds from cabbage outer leaves via the application of ultrasonic pretreatment. *Sep. Purif. Technol.* **144**:37–45.
- Provesi, J. G. and Amante, E. R. (2015). Carotenoids in pumpkin and impact of processing treatments and storage. In: *Processing and Impact on Active Components in Food*, pp. 71–80. Preedy, V. R., Ed., Elsevier, San Diego, CA.
- Puértolas, E., Cregenzán, O., Luengo, E., Álvarez, I. and Raso, J. (2013). Pulsed electric field-assisted extraction of anthocyanins from purple-fleshed potato. *Food Chem.* **136**:1330–1336.
- Qian, C., Decker, E. A., Xiao, H. and McClement, D. J. (2012). Physical and chemical stability of β -carotene-enriched nanoemulsions: Influence of pH, ionic strength, temperature, and emulsifier type. *Food Chem.* **132**:1221–1229.
- Rastogi, N. K. (2011). Opportunities and challenges in application of ultrasound in food processing. *Crit. Rev. Food Sci. Nutr.* **51**:705–722.
- Reshmi, S. K., Aravindhan, K. M. and Devi, P. S. (2012). The effect of light, temperature, pH on stability of betacyanin pigment in *basella alba* fruit. *Asian J. Pharm. Clin. Res.* **5**:107–110.

- Reynoso, R., Garcia, F. A., Morales, D. and de Mejia, E. G. (1997). Stability of betalain pigments from a cactacea fruit. *J. Agric. Food Chem.* **45**:2884–2889.
- Rodríguez-Saona, L. E., Giusti, M. M. and Wrolstad, R. E. (1999). Color and pigment stability of red radish and red-fleshed potato anthocyanins in juice model systems. *J. Food Sci.* **64**:451–456.
- Sharif, A., Saim, N., Jasmani, H. and Ahmad, W. Y. W. (2010). Effects of solvent and temperature on the extraction of colorant from onion (*Allium cepa*) skin using pressurized liquid extraction. *Asian J. Appl. Sci.* **3**:262–268.
- Silva, L. V., Nelson, D. L., Drummond, M. F. B., Dufossé, L. and Glória, M. B. A. (2005). Comparison of hydrodistillation methods for the deodorization of turmeric. *Food Res. Int.* **38**:1087–1096.
- Socaciu, C. (2008). Updated technologies for extracting and formulating food colorants. In: *Food Colorants: Chemical and Functional Properties*, pp. 303–322. Socaciu, C., Ed., CRC Press, Boca Raton, FL.
- Stamatopoulos, K., Katsoyannos, E., Chatzilazarou, A. and Konteles, S. J. (2012). Improvement of oleuropein extractability by optimising steam blanching process as pre-treatment of olive leaf extraction via response surface methodology. *Food Chem.* **133**:344–351.
- Steet, J. A. and Tong, C. H. (1996). Degradation kinetics of green color and chlorophylls in peas by colorimetry and HPLC. *J. Food Sci.* **61**:924–928.
- Stintzing, F. C. and Carle, R. (2008). Betalains in food: Occurrence, stability, and post-harvest modifications. In: *Food Colorants: Chemical and Functional Properties*, pp. 277–290. Socaciu, C., Ed., CRC Press, Boca Raton, FL.
- Takyi, E. E. K. (2001). Bioavailability of carotenoids from vegetables versus supplements. In: *Vegetables, Fruits and Herbs in Health Promotion*, pp. 19–34. Watson, R. R., Ed., CRC Press, Boca Raton, FL.
- Tan, J. B. L., Lim, Y. Y. and Lee, S. M. (2014). *Rhoeo spathacea* (Swartz) stearn leaves, a potential natural food colorant. *J. Funct. Foods.* **7**:443–451.
- Tao, Y. and Sun, D. W. (2015). Enhancement of food processes by ultrasound: A review. *Crit. Rev. Food Sci. Nutr.* **55**:570–594.
- Tao, Y., Wu, D., Zhang, Q. A. and Sun, D. W. (2014). Ultrasound-assisted extraction of phenolics from wine lees: Modeling, optimization and stability of extracts during storage. *Ultrason. Sonochem.* **21**:706–715.
- Tonucci, L. H. and von Elbe, J. H. (1992). Kinetics of the formation of zinc complexes of chlorophyll derivatives. *J. Agric. Food Chem.* **40**:2341–2344.
- Vági, E., Simándi, B., Daood, H. G., Deák, A. and Sawinsky, J. (2002). Recovery of pigments from *Origanum majorana* L. by extraction with supercritical carbon dioxide. *J. Agric. Food Chem.* **50**:2297–2301.
- Veggi, P. C., Martinez, J. and Meireles, M. A. A. (2013). Fundamentals of microwave extraction. In: *Microwave-Assisted Extraction for Bioactive Compounds: Theory and Practice*, pp. 15–52. Chemat, F. and Cravotto, G., Eds., Springer, New York, NY.
- Viña, S. Z., Olivera, D. F., Marani, C. M., Ferreyra, R. M., Mugridge, A., Chaves, A. R. and Mascheroni, R. H. (2007). Quality of Brussels sprouts (*Brassica oleracea* L. gemmifera DC) as affected by blanching method. *J. Food Eng.* **80**:218–225.
- von Elbe, J. H. and Schwartz, S. J. (1996). Colorants. In: *Food Chemistry*, 3rd ed., pp. 651–722. Fennema, O. R., Ed., Marcel Dekker, New York, NY.
- Wang, J. X., Xiao, H. X. and Li, G. K. (2008). Study of vacuum microwave-assisted extraction of polyphenolic compounds and pigment from Chinese herbs. *J. Chromatogr. A.* **1198–1199**:45–53.
- Wu, X. and Prior, R. L. (2005). Identification and characterization of anthocyanins by high-performance liquid chromatography-electrospray ionization-tandem mass spectrometry in common foods in the United States: Vegetables, nuts, and grains. *J. Agric. Food Chem.* **53**:3101–3113.
- Xiao, X., Song, W., Wang, J. and Li, G. (2012). Microwave-assisted extraction performed in low temperature and in vacuo for the extraction of labile compounds in food samples. *Anal. Chim. Acta.* **712**:85–93.
- Yang, Z. and Zhai, W. (2010). Optimization of microwave-assisted extraction of anthocyanins from purple corn (*Zea mays* L.) cob and identification with HPLC-MS. *Innovative Food Sci. Emerging Technol.* **11**:470–476.
- Yuhás Jr., S. A. (1995). Solvents. In: *Paint and Coating Testing Manual*, 4th ed., pp. 125–158. Koleske, J. V., Ed., ASTM, Ann Arbor, MI.
- Zou, T., Wu, H., Li, H., Jia, Q. and Song, G. (2013). Comparison of microwave-assisted and conventional extraction of mangiferin from mango (*Mangifera indica* L.) leaves. *J. Sep. Sci.* **36**:3457–3462.